

Irritable Bowel Syndrome

What is irritable bowel syndrome?

Irritable bowel syndrome (IBS) is a chronic (long-lasting) disorder of the large intestine (also called the colon or bowel). IBS is not a disease. It's a functional disorder, which means that the bowel doesn't work as it should. Although IBS can cause much distress, it does not damage the bowel and does not lead to life-threatening illness.

IBS is the most common intestinal disorder. It affects twice as many women as men and usually begins in early adult life.

How does it occur?

The cause of IBS is not fully understood. Most symptoms of IBS probably results from abnormal muscle contraction of the muscles in the lower part of the colon. Sometimes an abnormal contraction delays bowel movements, causing constipation. At other times it may lead to more rapid passage of bowel movements, causing diarrhea. In either case it usually causes abdominal cramps.

Some foods may trigger attacks. Symptoms may also be triggered by emotional stress or depression because muscles in the bowel are controlled by the nervous system.

What are the symptoms?

The most common symptoms include:

- cramping and pain in the abdomen, which may be severe
- constipation or diarrhea
- a lot of gas.

Other symptoms include:

- bloating
- a feeling of fullness in the rectum.
- Symptoms often occur after you have eaten a big meal or when you are under stress. Your symptoms may be relieved by having a bowel movement.

How is it diagnosed?

After taking your medical history, your health care provider will examine your abdomen and may do a rectal exam. There is no specific test for IBS. Depending on your history and exam, your provider may do the following tests to look for other possible causes of your symptoms:

- blood tests
- tests of bowel movement samples to check for blood and infection
- x-rays
- colonoscopy or sigmoidoscopy (procedures that allow your provider to see the inside of your colon with a thin, flexible, lighted tube)
- barium enema (a procedure in which a special liquid is passed into the colon through the rectum before x-rays are taken) to check the colon lining.

Your health care provider may ask you to try a milk-free diet to see if lactose intolerance (trouble digesting milk) may be causing your symptoms.

How is it treated?

Doctors have not yet found a cure for IBS. However, controlling the diet and emotional stress usually relieves the symptoms. Some medicines may also help.

- **Diet** Increasing the fiber in your diet often helps, although sometimes a decrease in fiber is needed. Try eating smaller meals more often. Avoid foods that cause gas, such as cabbage.
- **Food diary** Your health care provider may ask you to keep a food diary to see if eating a particular food, for example, milk or sausage, worsens your symptoms.
- **Stress** Your provider will help you identify things that cause stress in your life and will suggest ways to help you control them. Relaxation or biofeedback techniques may help you manage stress.
- **Medicines** Your provider may prescribe:
 - bulk-forming agents, such as bran or methylcellulose
 - antispasmodic drugs to slow contractions in the bowel and help with diarrhea and pain
 - antidepressants, which can help control chronic pain
 - one of two new serotonin-related medicines to help with diarrhea or constipation.

How long will the effects last?

Because IBS is a chronic disorder, you may have flare-ups of symptoms throughout your life. Although a cure hasn't been found yet, the disorder can usually be controlled. IBS will not progress to something worse.

How can I take care of myself?

- Follow your health care provider's recommendations.
- Learn stress-management techniques to reduce stress and anxiety in your life. Professional counseling may be helpful.
- Drink plenty of water.
- Do not drink alcohol, which can make symptoms of IBS worse.
- Select your foods carefully. If a food appears to bring on your symptoms, avoid it. However, don't eliminate a food just because it appears to cause symptoms one time. Be sure that a food produces symptoms several times before you give it up. You should try to keep many different foods in your diet because a varied diet provides better nutrition.
- Ask your health care provider if you should have a high-fiber diet, especially if you tend to be constipated. High-fiber foods may cause gas and bloating, but usually these symptoms lessen as the digestive tract gets used to the increased fiber. Some high-fiber foods include:
 - whole-grain breads and cereals
 - fruits, especially apricots, blackberries, coconut, dates, figs, kiwi, peaches, pears, pineapple, prunes, raspberries, and strawberries
 - nuts, especially almonds, pistachios, and walnuts
 - vegetables, particularly Brussels sprouts, corn and popcorn, broccoli, and parsley
 - beans and lentils.
- Ask your health care provider if you should use a nonprescription fiber supplement.
- Eat smaller meals more often. For example, eat 6 small meals a day rather than 3 large ones.
- See your health care provider if your symptoms are getting worse or you are having them more often.

Published by McKesson Health Solutions LLC. The information is intended to inform and educate and is not a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional. Developed by McKesson Health Solutions LLC. Copyright © 2003 McKesson Health Solutions LLC. Copyright © Clinical Reference Systems 2004 **Adult Health Advisor** Copyright © 2004 Elsevier Inc. All rights reserved.

Special Instructions: