

Upper Back Pain

What is upper back pain?

Your upper back is also called your thoracic back, the part of the back where the ribs attach. Upper back pain is pain between your neck and your low back.

How does it occur?

The bones in your back are called vertebrae. Back pain is usually caused when ligaments or muscles attaching to the vertebrae are injured. Upper back pain can come from a twisting motion, poor posture, overuse, or an injury. It is very common for someone to injure their upper back when carrying objects, throwing, bending or twisting. Sitting at a desk for a prolonged time can cause upper back muscles to tighten and become stiff. Upper back pain can come even come from vigorous coughing or sneezing.

Sometimes upper back pain is caused by scoliosis, a curve in the spine that has developed during the adolescent growth period. In scoliosis there is usually an imbalance of the muscles of the upper back.

What are the symptoms?

- Symptoms of upper back pain may include:
- pain in the upper back
- muscle spasms
- pain when you take a deep breath
- pain when your back is touched or when you move
- pain when you move your shoulders or bend your neck forward.

How is it diagnosed?

Your provider will take your history, review your symptoms and examine your back.

How is it treated?

The early stages of back pain with muscle spasms should be treated with ice packs for 20 to 30 minutes every 4 to 6 hours for the first 2 to 3 days. You may use a frozen gel pack, crushed ice, or a bag of frozen peas. After you have iced for 2 to 3 days, you may start to use moist heat to help loosen up stiff muscles.

Your provider may prescribe an anti-inflammatory medication, muscle relaxants, or other medication. Massage to the inflamed muscles will help. Your provider will recommend exercises to help your back.

When can I return to my sport or activity?

The goal of rehabilitation is to return you to your sport or activity as soon as is safely possible. If you return too soon you may worsen your injury, which could lead to permanent damage. Everyone recovers from injury at a different rate. Return to your sport or activity will be determined by how soon your back recovers, not by how many days or weeks it has been since your injury occurred. In general, the longer you have symptoms before you start treatment, the longer it will take to get better.

It is important that you have fully recovered from your upper back pain before you return to your sport or strenuous activity. You must be able to have the same range of motion that you had before the injury. You must be able to run, lift, jump and twist without pain.

What can I do to prevent upper back pain?

Be sure that you have warmed up and have done proper stretching exercises before your activity. Try not to twist when you are lifting heavy objects. If you are at a desk for a long period of time be sure to take frequent breaks to stretch you back.

Published by McKesson Health Solutions LLC.

This content is reviewed periodically and is subject to change as new health information becomes available. The information is intended to inform and educate and is not a replacement for medical evaluation, advice, diagnosis or treatment by a healthcare professional.

Written by Pierre Rouzier, MD.

Copyright © 2003 McKesson Health Solutions LLC. All rights reserved.

Special Instructions:

Copyright © Clinical Reference Systems 2004

Copyright © 2004 Elsevier Inc. All rights reserved.

www.mdconsult.com